
mCME QUESTIONNAIRE

Name: Country: City: P.O.Box:

Email address: Tel: Mob: Membership number:

Please email back to events@cappmea.com

Long-term clinical success in the management of compromised intertooth spaces utilizing small-diameter implants

by Paul S. Petrunaro, DDS, MS

1. What is the solution to insufficient intertooth spacing between natural teeth?
 - a) removable partial denture
 - b) resin bonded bridge
 - c) fixed bridge
 - d) all of the above

2. Wallace, Misch and Salama stated that for the normal two-piece implant, the implant should be placed:
 - a) at least 1.5 mm
 - b) at least 1.6 mm
 - c) at least 1.7 mm

3. Small-diameter, or mini, implants were developed:
 - a) less than 20 years ago
 - b) more than 20 years ago

4. Implant are available from 1.8 mm diameter to 2.8 mm diameter:
 - a) True
 - b) False

5. Was the use of small-diameter or mini implants expanded into immediate conversion of full dentures into implant-supported dentures?
 - a) True
 - b) False

6. The implants that are available from 1.8 mm diameter to 2.8 mm diameter, offer a fixed permanent tooth replacement option for patients who otherwise would not be able to have implants placed and restored.
 - a) True
 - b) False

mCME QUESTIONNAIRE

Name: Country: City: P.O.Box:

Email address: Tel: Mob: Membership number:

Please email back to events@cappmea.com

7. What is the smallest small-diameter implant from most dental implants manufacturers?
 - a) 3.0 mm
 - b) 2.8 mm
 - c) 2.5 mm

8. Using dental implants to replace the natural tooth system in the esthetic zone has also seen an increase in restorative treatment plans
 - a) True
 - b) False

9. Patients seeking tooth replacement for partial or totally edentulous situations have been able to enjoy natural appearing and functioning prostheses that are fixed, stable and, in some cases, so natural it's difficult to ascertain a dental implant restoration from a tooth restoration
 - a) True
 - b) False

10. To properly form an ovate pontic type emergence profile in the soft tissue, which is required for a fixed bridge to have a natural clinical appearance, consideration must be given to the intertooth edentulous space
 - a) True
 - b) False

Evaluation Form

This represented new knowledge: None Some Moderate New

Did the article meet your learning objectives? Yes No

Was the article relevant to the topic? Yes No

Did the article meet your expectations? Yes No

In order to serve you better, we would like to know your preferred topics for the next issues:

.....